

# Fresh-cut Vegetables

M. Margaret Barth<sup>1</sup>, Hong Zhuang<sup>1</sup> and Mikal E. Saltveit<sup>2</sup>

<sup>1</sup>Redi-Cut Foods, Inc., Franklin Park, IL

<sup>2</sup>Mann Lab, Dept. of Vegetable Crops, University of California, Davis, CA

**Introduction:** Wounds inflicted during the preparation of fresh-cut vegetables promote many physical and physiological changes that hasten loss of product quality (Brecht, 1995; Saltveit, 1997). Foremost among these, are the removal of the protective epidermal layer and/or exposure of internal cells. These changes not only facilitate water loss, but also provides an easy entry for microbial pathogens and chemical contaminants. Packaging and/or application of edible films can lessen water loss by maintaining a high RH near the cut surface, and providing a physical barrier that protects the product from contamination.

Water loss and collapse of injured cells at the cut surface can alter the appearance of the fresh-cut product. As the cut surface loses water, adhering cellular debris may impart a white blush to the surface that masks varietal color, eg., white blush on 'baby' carrots decreases the intensity of the underlying orange color. Differential dehydration of exposed cortex and vascular tissue may produce an uneven surface, eg., vascular strands projecting a few millimeters from the cut end of celery petioles. Consumers associate both of these surface changes with the loss of freshness.

Physiological processes devoted to wound repair can be beneficial (eg., curing of potatoes) or detrimental to quality retention. Physiological changes following wounding include increased respiration and ethylene production, promotion of the ripening of climacteric fruit vegetables (eg., melons, tomatoes), and enhanced synthesis and accumulation of phenolic compounds that contribute to tissue browning. These changes are managed by the use of low temperatures, creation of reduced O<sub>2</sub> and/or elevated CO<sub>2</sub> atmospheres, and the application of inhibitors of specific chemical reactions or metabolic pathways.

Wound enhanced respiration and ethylene production can deplete carbohydrate storage reserves and stimulate tissue softening associated with fruit ripening (eg., melons, squash, tomatoes) or chlorophyll loss associated with leaf or tissue senescence (eg., spinach, broccoli). In many tissues, wounding produces a signal that induces the increased synthesis and accumulation of phenolic compounds. Tissue browning can result from the oxidation and/or polymerization of accumulated phenolics compounds. Lignification and toughening of tissue are others way in which the metabolism of phenolic compounds can reduce quality.

Sanitation, implementation of Good Agricultural Programs (GAP's) in the field, use of Good Manufacturing Programs (GMP's) and Hazard Analysis and Critical Control Point (HACCP) monitoring plans during all steps of processing, and proper temperature control, are all needed to insure that an initial low microbial load on fresh-cut vegetables is maintained during marketing. The current industry standard washing and disinfection procedures used on whole produce are rather ineffective with fresh-cut vegetables once they have been contaminated with microbial pathogens. Overall microbial load relative to spoilage organisms usually remain below levels of concern as long as visual quality remains acceptable. However, postharvest treatments that maintain visual quality under abusive temperatures may allow microbial loads to reach dangerous levels before quality is reduced below acceptable levels.

Many vegetables are sensitive to non-freezing temperatures below around 50 °F (10 °C), and suffer physiological damage if held at these chilling temperatures beyond a specific period of time. These vegetables include jicama, pepper, sweet potato, tomato and zucchini. In contrast to the whole commodity that should not be chilled, the increased susceptibility of fresh-cut products to spoilage often means that the product has a longer market life at 32 °F (0 °C) than at higher, non-chilling temperatures. Development of chilling injury symptoms is less pronounced in riper tissue, and takes some time to develop. Since most fresh-cut vegetables are consumed soon after purchase and fruit vegetables are fully ripe when processed, chilling injury symptoms usually do not develop to a significant extent because the rate of deterioration due to chilling is slower than the rate of spoilage that is inhibited by low temperatures. The two most important

fresh-cut commodities, lettuce and carrots, are not chilling sensitive and should be stored as close to 32 °F (0 °C) as possible without freezing, ie., at 34 to 38 °F (1 to 3 °C) in a commercial setting.

The imposed physical damage done to the product during preparation, and its increased vulnerability to deleterious internal and external changes, requires that fresh-cut vegetables be handled with a greater degree of care than the whole product. Storage requirements for intact produce may be inadequate to handle increased physiological activity and susceptibility to water loss encountered with fresh-cut vegetables.

The highest quality that is available (usually USDA No. 1) and that is compatible with an economic return should be used for processing. Although poor quality pieces of fresh-cut product can be discarded to upgrade quality (eg., culling discolored pieces of cut lettuce leaves or broccoli florets), the associated cost usually adversely impacts quality and yields, and greatly outweighs the differential cost of using higher quality raw material.

The remainder of this chapter provides information on the commercial packaging and storage of a number of fresh-cut vegetables.

### **BEETS, Red (grated, cubed, whole peeled)**

Fresh-cut beets should be stored at 34 to 38 °F (1 to 3 °C) before and after processing. Respiration is slightly reduced during storage in 5% O<sub>2</sub> + 5% CO<sub>2</sub> at 5 °C (41 °F).

#### **Respiration Rates:**

Temperature	Whole Peeled	Cubed (1 x 1 cm)	Grated (2 mm)
	(mg CO <sub>2</sub> kg <sup>-1</sup> h <sup>-1</sup> )		
2 °C	4	10	12
5 °C	6	12	16
10 °C	19	27	38
23 °C	54	117	162 to 207

To get mL kg<sup>-1</sup> h<sup>-1</sup>, divide the mg kg<sup>-1</sup> h<sup>-1</sup> rate by 2.0 at 0 °C (32 °F), 1.9 at 10 °C (50 °F), and 1.8 at 20 °C (68 °F). To calculate heat production, multiply mg kg<sup>-1</sup> h<sup>-1</sup> by 220 to get BTU per ton per day or by 61 to get kcal per metric ton per day. Data from Gorny, 1997.

### **BROCCOLI (Florets)**

Fresh-cut broccoli florets should be tight, firm, turgid and dark green without blooming buds. There should be no sulfur odor or discoloration along the stems and cut ends. The core temperature of raw material should be < 1.5 °C (35 °F). Raw and processed material should be stored at 34 to 38 °F (1 to 3 °C) to assure quality and reduce potential for freezing of product during handling, distribution and storage. Yellowing is a common problem caused by either chlorophyll loss or blooming of the buds. The cut surface and damaged floret stems can turn black during storage. Development of off-odors can be a major concern when MAP is used. Temperature abuse promotes soft rot and mold growth.

Whole broccoli heads are hand-cut into florets that are > 1 in (2.5 cm) to < 2 in (5 cm) long. They are washed in water containing up to 200 ppm total chlorine to wash residual material from the florets as well as reduce aerobic plate counts. Repeated daily washing did not maintain lower microbial counts for > 2 days.

The benefit of CA (5% O<sub>2</sub> + 4% CO<sub>2</sub>) may be marginal for storage up to 14 days at 32 to 41 °F (0 to 5 °C) compared to air. Lowering O<sub>2</sub> to 0.25% or increasing CO<sub>2</sub> to 10% at 32 to 41 °F (0 to 5 °C) reduces respiration by about 50%. Use of appropriate polymeric film in MAP maintained green color at 32 to 41 °F (0 to 5 °C) for more than 21 days (Cabezas and Richardson, 1997). Severe off-odors and discoloration at cut ends can develop during MAP storage at 10% CO<sub>2</sub> + 2.5% or less O<sub>2</sub> (Makhlouf et al., 1989). Perforated and micro-perforated polymeric packages reduce off-odor (Izumi et al., 1996a). Reducing ethylene below 1 to 10 µL L<sup>-1</sup> did not significantly reduce color loss at 34 °F (1 °C), but did have an affect at higher

temperatures.

No food-borne disease outbreak has been reported associated with fresh-cut broccoli although the aerobic population is usually high (> 100,000 cfu/g fresh weight). Florets stored in 5% O<sub>2</sub> + 8% CO<sub>2</sub> at 46 °F (8 °C) have lower aerobic plate counts, total coliform plate counts, and yeast/mold plate counts compared to storage in air.

#### **Respiration Rates: fresh-cut florets**

Temperature	mg CO <sub>2</sub> kg <sup>-1</sup> h <sup>-1</sup>
0 °C	26
5 °C	44
10 °C	78

To get mL kg<sup>-1</sup> h<sup>-1</sup>, divide the mg kg<sup>-1</sup> h<sup>-1</sup> rate by 2.0 at 0 °C (32 °F), 1.9 at 10 °C (50 °F), and 1.8 at 20 °C (68 °F). To calculate heat production, multiply mg kg<sup>-1</sup> h<sup>-1</sup> by 220 to get BTU per ton per day or by 61 to get kcal per metric ton per day. Data from Watada et al. (1996) and Izumi et al. (1996a).

#### **CABBAGE, Green and Red (shredded, diced)**

Fresh-cut green cabbage should be light green with a moderate pungent flavor and no sulfur aroma. Raw and processed material should have typical cabbage flavor with no off notes, and should be stored at 34 to 38 °F (1 to 3 °C) to assure quality and reduce potential for freezing of product during handling, distribution and storage. Fresh-cut cabbage includes diced and shredded product with cut size varying from ¼ in (0.63 cm) to 3/8 in (0.95 cm). Heads are trimmed to remove wrapper leaves and cored, cut, and washed using chlorinated water (100 ppm of total chlorine) for about 1 min before being spun-dried and packaged. For coleslaw, cut cabbage (shredded or diced) can be pre-blended with carrots or plain-packaged.

A CA of 5 to 7.5% O<sub>2</sub> + 15% CO<sub>2</sub> is recommended (Hiroaki et al., 1993). Lowering O<sub>2</sub> below 5% caused rapid proliferation of fermentative bacteria and off odors within 6 days at 5°C. The fermentation induction point of coleslaw mixes in low density polyethylene bags varied with the cabbage to carrot ratio, and temperature. At 41 °F (5 °C), a 70:30 mix went anaerobic at 1.8% O<sub>2</sub>, while 3% O<sub>2</sub> was the limit for aerobic respiration at 10°C.

Off-odors formation in cut cabbage bags with low OTR film (< 3,000 mL/m<sup>2</sup> atm<sup>-1</sup> day<sup>-1</sup>) and discoloration of cabbage leaf packed in high OTR film (> 12,000 mL/m<sup>2</sup> atm<sup>-1</sup> day<sup>-1</sup>) bags are the major quality deterioration during storage (Pirovani et al., 1997). At 52 °F (11 °C), *Listeria* spp grew faster in fresh-cut cabbage packages which have an atmosphere of < 1.8% O<sub>2</sub> + > 20% CO<sub>2</sub> than in air (Omary et al., 1993).

#### **Respiration Rates: shredded**

Temperature	3/8 in (9.6 mm)	1/4 in (6.4 mm)
	(mg CO <sub>2</sub> kg <sup>-1</sup> h <sup>-1</sup> )	
2 °C	16 to 18	18 to 24
5 °C	22 to 33	25 to 39
10 °C	42 to 48	51 to 57
23 °C	117 to 153	153 to 171

To get mL kg<sup>-1</sup> h<sup>-1</sup>, divide the mg kg<sup>-1</sup> h<sup>-1</sup> rate by 2.0 at 0 °C (32 °F), 1.9 at 10 °C (50 °F), and 1.8 at 20 °C (68 °F). To calculate heat production, multiply mg kg<sup>-1</sup> h<sup>-1</sup> by 220 to get BTU per ton per day or by 61 to get kcal per metric ton per day. Data from Watada et al. 1996

**CABBAGE, CHINESE (sliced, sticks, shredded)**

Fresh-cut Chinese cabbage should be handled like cabbage and stored at 34 to 38 °F (1 to 3 °C). The recommended CA is 5% O<sub>2</sub> + 5% CO<sub>2</sub>; benefits are moderate compared to air storage at 32 °F (0 °C).

**CARROTS (diced, shredded, sticks, peeled, grated, sliced, cubes)**

Fresh-cut carrots are orange without a white brush or slimy surface. Raw and processed product should be kept at 34 to 38 °F (1 to 3 °C) to assure quality and reduce potential for freezing of product during handling, distribution and storage. Most quality loss in cut carrots results from either formation of white blush, or off-odor and slimy surface generated by bacteria (Carlin et al., 1989).

Fresh-cut carrots include whole peeled (baby), sticks, sliced, shredded, grated and diced. Whole carrots are washed with water to remove undesirable field material. The stems and tips are excised and the trimmed carrots are peeled, cut, and washed in 100 µL L<sup>-1</sup> NaOCl for less than 1 min. Washed carrot cuts are centrifuged to remove excess water and packaged in plastic bags.

Dehydration of surface debris on cut and peeled carrots imparts a whitish translucent appearance to the surface (Tatsumi et al., 1991; Cisneros-Zevallo et al., 1995), which is undesirable because consumers associate it with the loss of freshness (Bolin and Huxsoll, 1991). Application of edible coating (Howard and Dewi, 1995, 1996, Li and Barth, 1998), such as sodium caseinate-stearic acid (Avena-Bustillos et al., 1994) or heating and raising the pH (Bolin and Huxsoll, 1991) may be helpful in reducing white blush. Treatments that modifying the water retaining capacity of the cut surface also prevent white blush development (Cisneros-Zevallos et al., 1997).

Fresh-cut carrots derive slight benefit from 2 to 5% O<sub>2</sub> + 15 to 20% CO<sub>2</sub> atmospheres (Izumi et al., 1996b). Lower O<sub>2</sub> or increased CO<sub>2</sub> levels promoted slimy appearance, increased lactic acid bacteria growth, and accelerated microbial decay and excessive alcohol production. Grated carrots retain good quality for up to 10 days at 36 to 52 °F (2 to 10 °C) in MAP with high O<sub>2</sub> permeability films of 10 to 20 L m<sup>2</sup> atm<sup>-1</sup> day<sup>-1</sup> at 25°C (Carlin et al., 1990). Low permeability films of 950 mL m<sup>2</sup> atm<sup>-1</sup> day<sup>-1</sup> resulted in low O<sub>2</sub> damage.

The quality and headspace composition of sliced carrots (0.5 cm) treated with 1% ascorbic acid before packing in MAP was unaffected during 14 days at 40 °F (4 °C) (Galetti et al. 1997). Growth of aerobic mesophilic bacteria on sticks was suppressed by a 0.5% O<sub>2</sub> + 10% CO<sub>2</sub> atmosphere at both 32 and 41 °F (0 and 5 °C); but total microbial count on slices and shreds was unaffected.

The mean microbial population after 9 days storage was much lower (1,300 CFU g<sup>-1</sup>) for irradiated shredded carrots (0.5 kGy) than for non-irradiated, chlorinated controls (87,000 CFU g<sup>-1</sup>) (Hagenmaier and Baker, 1998). Ethanol and O<sub>2</sub> content of the headspace were not affected.

**Respiration Rates:**

Temperatures	Whole-peeled	Sliced (mg CO <sub>2</sub> kg <sup>-1</sup> h <sup>-1</sup> )	Sticks	Shredded
0 °C	---	5	11	15
5 °C	9 to 12	13	19	24
10 °C	17 to 21	25	42	46
23 °C	54	72 to 81	---	108 to 126

To get mL kg<sup>-1</sup> h<sup>-1</sup>, divide the mg kg<sup>-1</sup> h<sup>-1</sup> rate by 2.0 at 0 °C (32 °F), 1.9 at 10 °C (50 °F), and 1.8 at 20 °C (68 °F). To calculate heat production, multiply mg kg<sup>-1</sup> h<sup>-1</sup> by 220 to get BTU per ton per day or by 61 to get kcal per metric ton per day. Data from Gorny 1997.

**CELERY (diced, sliced, sticks)**

Fresh-cut celery petioles have no leaves or brown spots, and should not have cracked, flared or white-blush ends. Raw material should arrive at < 38 °F (7 °C) and be stored at 36 to 40 °F (2 to 4 °C) before processing,

and at 34 to 38 °F (1 to 3 °C) after processing, to assure quality and reduce potential for freezing of product during handling, distribution and storage. Atmospheres of 5% O<sub>2</sub> + 4% CO<sub>2</sub> were slightly beneficial. Water loss is the main factor affecting eating quality (Garipey et al., 1986). Small amounts of water loss (2.5 to 5%) can result in litheness, flaccidity, shriveling and wrinkling. Moisture loss can be reduced 75% by the application of an edible coating (Avena-Bustillos et al., 1997). Differential dehydration of exposed cortex and vascular tissue caused the stronger vascular strands to project a few millimeters from the cut end of celery petioles. Development of pithiness is associated with water stress during growth and wounding during processing (Saltveit and Mangrich, 1996). Heat-shock treatments prevented tissue browning without adversely affecting other quality parameters (Loaiza-Velarde et al., 2002).

Sliced celery can only be maintained for 6 days at 40 °F (4.4°C) without notable quality change (Johnson et al., 1974). Major pathogens are *Pseudomonas fluorescens* and *P. Marginalis*, which cause water soaking, soft rot, and discoloration (Robbs et al., 1996). Low-dose irradiation (1.0 kGy) delayed microbial spoilage in fresh, diced celery stored at 40 °F (4 °C) for 3 weeks without affecting sensory attributes. In contrast, blanching and chlorine rinsing had little effect on microbial count. Sensory results indicated the 1 kGy sample was preferred over all other treatments in terms of taste, texture and color.

### **GARLIC (peeled)**

Quality problems with peeled garlic include sprouting and discoloration of areas damaged during peeling. Garlic cloves can be peeled manually or mechanically with compressed air. Acid treatments and edible coatings can improve preservation. Excellent visual quality was maintained for 21, 16, 12 or 8 days of storage in air at 32, 41, 50, or 59 °F (0, 5, 10, or 15 °C), respectively. Decay was not observed at 32 or 41 °F (0 and 5 °C), and was slight at 50 and 59 °F (10 and 15 °C). At 41 and 50 °F (5 and 10 °C), a 1% O<sub>2</sub> + 10% CO<sub>2</sub> CA gave the best quality (lack of decay and least discoloration). Respiration rates in manually peeled garlic were 50% higher than unpeeled cloves. Mechanically-peeled garlic has rates of respiration 5 to 10% higher at 41 °F (5 °C) and 20 to 30% higher at 50 °F (10 °C), than hand-peeled cloves.

### **Respiration Rate: peeled**

Temperature	mg CO <sub>2</sub> kg <sup>-1</sup> h <sup>-1</sup>
5 °C	35
10 °C	57

To get mL kg<sup>-1</sup> h<sup>-1</sup>, divide the mg kg<sup>-1</sup> h<sup>-1</sup> rate by 2.0 at 0 °C (32 °F), 1.9 at 10 °C (50 °F), and 1.8 at 20 °C (68 °F). To calculate heat production, multiply mg kg<sup>-1</sup> h<sup>-1</sup> by 220 to get BTU per ton per day or by 61 to get kcal per metric ton per day. Data from Burlo-Carbonell et al., 2000.

### **JICAMA (sticks, cubes)**

Fresh-cut jicama should be white with a crisp texture. Raw material should be received at < 44 °F (7 °C), and processed product stored at 34 to 38 °F (1 to 3 °C) to assure quality and reduce potential for freezing of product during handling, distribution and storage. Tuber size does not affect visual quality or color. Browning can be a significant cause of quality loss after 9 days at 41 °F (5 °C). Increased storage temperature results in accelerated discoloration. The best treatment to control browning is a 2-min dip in water at 122 °F (50 °C) and then storage at 36 °F (2 °C).

Quality loss during storage at 0 or 5°C in air is also attributable to yeast and bacterial growth. A 10% CO<sub>2</sub> atmosphere retards brown discoloration, maintains crisp texture, and delays microbial growth. However, acetaldehyde and ethanol concentrations are higher after 8 days at 41 °F (5 °C) in 10% CO<sub>2</sub> atmospheres + 3% O<sub>2</sub> or air. Atmospheres of 20% CO<sub>2</sub> can damage jicama if stored longer than 8 days.

### **KALE (shredded)**

Fresh-cut kale should be dark green, crisp and without yellow or discolored leaf pieces. Raw material should be received at < 45 °F (7 °C) and stored at 34 to 38 °F (1 to 3 °C) after processing to assure quality and reduce potential for freezing of product during handling, distribution and storage. Washing shredded kale with 0.1% NaOCl results in a 10-fold reduction in aerobic plate count compared to unwashed products.

Fresh-cut kale in perforated ( $5.96 \pm 0.35 \text{ mm}^2$  diameter holes with  $6.68 \text{ cm}^2$  total surface area per sack) polyethylene bags (38  $\mu\text{m}$  thick) become unacceptable due to yellowing and microbial spoilage after 3 days at 39 °F (4 °C). Desiccation, yellowing and spoilage were unacceptable after 1 or 2 days at 68 or 50 °F (20 or 10 °C), respectively (Beaulieu et al., 1997).

### **LEEK (sliced)**

Raw material should be received at, and processed leeks stored at, 34 to 38 °F (1 to 3 °C) to assure quality and reduce potential for freezing of product during handling, distribution and storage. A CA of 5% O<sub>2</sub> + 5% CO<sub>2</sub> is moderately beneficial, and slightly reduces respiration at 41 °F (5 °C). A non-perforated film led to unacceptable material after a few days at 39 °F (4 °C) with high CO<sub>2</sub>, ethanol and acetaldehyde concentrations (Keteleer et al., 1993).

### **Respiration Rates:**

Temperature	Whole Leaf (mg CO <sub>2</sub> kg <sup>-1</sup> h <sup>-1</sup> )	Rings (2 mm thick) (mg CO <sub>2</sub> kg <sup>-1</sup> h <sup>-1</sup> )
2 °C	24	32
5 °C	29	49
10 °C	38	57 to 67
23°C	117	252 to 288

To get mL kg<sup>-1</sup> h<sup>-1</sup>, divide the mg kg<sup>-1</sup> h<sup>-1</sup> rate by 2.0 at 0 °C (32 °F), 1.9 at 10 °C (50 °F), and 1.8 at 20 °C (68 °F). To calculate heat production, multiply mg kg<sup>-1</sup> h<sup>-1</sup> by 220 to get BTU per ton per day or by 61 to get kcal per metric ton per day. Data from Gorny, 1997.

### **LETTUCE, Butterhead, Crisphead, Green leaf, Iceberg, Romaine; (chopped, shredded, whole leaf)**

The quality of the fresh-cut product is very dependent on the initial raw product quality. The raw product should be the best quality available, and should be < 36 °F (2 °C) when received as whole or cored heads.

It should be stored at 34 to 38 °F (1 to 3 °C) before and after processing to assure quality and reduce potential for freezing of product during handling, distribution and storage.

Browning (enzymatic) of the cut edges is the main defect of fresh-cut lettuce pieces during storage. The best way to prevent the discoloration is to reduce O<sub>2</sub> to < 3%. Antioxidant treatment and heat-shock can inhibit discoloration, but these treatments may result in loss of visual quality. Unless stressed in the field, lettuce has low levels of phenolic compounds and tissue browning occurs after the induction of synthesis and accumulation of phenolics by wounding. A 90 sec heat-shock at 104 °F (45 °C) prevented wound-induced browning of Iceberg (Loaiza-Velarde et al., 1997; Loaiza-Velarde and Saltveit, 2001) and Romaine (Saltveit, 2000) lettuce.

### ***Butterhead lettuce***

Browning of the cut edges of butterhead lettuce is reduced by modified atmospheres rapidly created by flushing with N<sub>2</sub> to get 1 to 3% residual O<sub>2</sub>, with CO<sub>2</sub> levels of 5 to 10%. Tissue in atmospheres of < 1% O<sub>2</sub> + > 10% CO<sub>2</sub> develop CO<sub>2</sub> injury or brown stain (Stewart and Ceponis, 1968). The incidence of brown stain increased from 16 to 38 to 81% as CO<sub>2</sub> increased from 2.5 to 5 to 10%, respectively, and was enhanced

when combined with low O<sub>2</sub> atmospheres (Steward and Uota, 1971).

The physiological behavior of butterhead lettuce, eg., sensitivity to CO<sub>2</sub> injury, is modified by cultural practices such as irrigation, climate and fertilization (Sorensen et al., 1994; Poulsen et al., 1994, 1995). Susceptibility to brown stain varies among cultivars and position of leaves within the head (Krahn, 1977).

Shelf-life of butterhead lettuce was negatively correlated with respiration rate and susceptibility to CO<sub>2</sub>. High O<sub>2</sub> and low CO<sub>2</sub> enhanced enzymatic browning. Off odors and flavors developed (ethanol, acetaldehyde and ethyl acetate aroma), crispness was lost, and the tissue became translucent when fresh-cut lettuce was stored in < 0.5% O<sub>2</sub> and/or > 15% CO<sub>2</sub>.

Necrosis of butterhead lettuce may be associated with microbial growth, such as pectolytic *Pseudomonas* (Nguyen-the and Prunier, 1989) and other epiphytic bacteria.

### **Respiration Rate: chopped**

Temperature	mg CO <sub>2</sub> kg <sup>-1</sup> h <sup>-1</sup>
2 °C	12 to 14
4.5 °C	20 to 25
10 °C	38 to 48

To get mL kg<sup>-1</sup> h<sup>-1</sup>, divide the mg kg<sup>-1</sup> h<sup>-1</sup> rate by 2.0 at 0 °C (32 °F), 1.9 at 10 °C (50 °F), and 1.8 at 20 °C (68 °F). To calculate heat production, multiply mg kg<sup>-1</sup> h<sup>-1</sup> by 220 to get BTU per ton per day or by 61 to get kcal per metric ton per day. Data from Gorny, 1997.

### ***Greenleaf lettuce***

A 0.5 to 3% O<sub>2</sub> + 5 to 10% CO<sub>2</sub> CA reduced cut edge browning of green leaf lettuce (Lopez-Galvez et al., 1996).

### ***Iceberg lettuce***

The raw product should have < 6% pink rib discoloration, while russet spotting, hooked core and tipburn should be < 1%. Moisture content of lettuce is extremely important for maintaining texture and reducing microbial growth. Excess moisture can be removed in a forced-air tunnel or by spinning. Lettuce that was air dried for 10 to 15 min or spin dried for 30 min had less browning during storage than control samples. Consumer sensory panelists indicated that after 4 days of storage, only 15% would purchase control lettuce while 40% would purchase lettuce spin or air dried for 15 min. Slicing methods can influence ascorbic acid retention in with the order: manual tearing > manual slicing > machine slicing.

In clamshell trays, market-life of fresh-cut lettuce was < 24 h at 73 °F (23 °C), > 24 h at 50 °F (10 °C), and > 48 h at 36 °F (2 °C) (Zhuang and Barth, 1999). Fresh-cut lettuce packages commonly have < 1% O<sub>2</sub> to effectively slow browning and > 10% CO<sub>2</sub> to inhibit microbial growth (Gorny, 1997). An atmosphere of 0.5 to 3% O<sub>2</sub> + 10 to 15% CO<sub>2</sub> reduces cut edge browning, retains visual quality and reduces Psychrophilic bacterial counts. Flushing with 100% N<sub>2</sub> increases retention over MAP. On the basis of ethanol appearance in the headspace, the fermentation induction point was 0.3 to 0.4% O<sub>2</sub> at 41 °F (5 °C) and 0.6 to 0.8% at 50 °F (10 °C). Acetaldehyde and ethyl acetate were detected when O<sub>2</sub> was < 0.4% at 41 to 68 °F (5 to 20 °C). The extent of browning increased with O<sub>2</sub> level, but not with CO<sub>2</sub> level (Smyth-Anne et al., 1998). Fresh-cut Iceberg should be stored at 34 to 38 °F (1 to 3 °C) before and after processing to assure quality and reduce potential for freezing during handling, distribution and storage.

Shelf-life of fresh-cut lettuce at 41 °F (5 °C) was 6 days in air, 8 days in 3% O<sub>2</sub>, and 12 days in 0.2% O<sub>2</sub> (Peiser et al., 1997). Increasing CO<sub>2</sub> enhances the beneficial effect of low O<sub>2</sub>. A combination of 0.2% O<sub>2</sub> with 7 to 15% CO<sub>2</sub> gave 16 day shelf-life. A 14 day shelf-life was obtained when shredded Iceberg lettuce was stored in 1 to 3% O<sub>2</sub> + 5 to 6% CO<sub>2</sub> at 41 °F (5 °C). However, CO<sub>2</sub> > 15% causes brown stain.

Reduced O<sub>2</sub> and/or elevated CO<sub>2</sub> may induce fermentation and off-odor (fermentative volatiles, ethanol and acetaldehyde) formation in the packages (McDonald et al., 1990; Peiser et al., 1997; Mateos et al., 1993).

Although gram-negative bacteria are numerically dominant, a large yeast population may also be found. Species of *Pseudomonas*, *Erwinia*, and *Serratia* were the most frequently isolated bacteria. *Cryptococcus*, *Pichia*, *Torulasporea* and *Trichosporon* spp. were the most common yeasts. Beuchat and Brackett (1990) studied the effects of shredding, chlorine treatment and MAP on survival and growth of *Listeria monocytogenes*, mesophilic aerobes, psychrotrophs, yeasts and molds. They found that no significant changes in populations of *L. monocytogenes* were detected during the first 8 days of incubation at 41 °F (5 °C); there was a significant increase between 8 and 15 days. Significant increases occurred within 3 days when lettuce was stored at 50 °F (10 °C). Chlorine treatment, MAP of 3% O<sub>2</sub>, and shredding did not influence growth of *L. monocytogenes*.

#### **Respiration Rate: shredded**

Temperature	mg CO <sub>2</sub> kg <sup>-1</sup> h <sup>-1</sup>
2.5 °C	12.0 to 15.0
5 °C	15.6 to 27.3
7.5 °C	23.1 to 32.7
10 °C	30.4 to 39.9

To get mL kg<sup>-1</sup> h<sup>-1</sup>, divide the mg kg<sup>-1</sup> h<sup>-1</sup> rate by 2.0 at 0 °C (32 °F), 1.9 at 10 °C (50 °F), and 1.8 at 20 °C (68 °F). To calculate heat production, multiply mg kg<sup>-1</sup> h<sup>-1</sup> by 220 to get BTU per ton per day or by 61 to get kcal per metric ton per day. Data from Gorny, 1997.

#### ***Lollo Rosso (Red) lettuce***

The beneficial storage atmosphere is 0.5 to 3% O<sub>2</sub> + 5 to 10% CO<sub>2</sub>. MAP storage (2 to 3% O<sub>2</sub> + 12 to 14% CO<sub>2</sub>) was useful in preventing brown discoloration. However, MAP storage was not beneficial for preserving quality of red tissues and, in fact, their overall visual quality, texture, aroma and macroscopic breakdown under MAP conditions were worse than those of air-stored tissues (Gil-Maria et al., 1998). Castaner et al. (1997) found that after 14 days of storage at 41 °F (5 °C), the overall visual quality of fresh-cut white and green tissue of Lollo Rosso (red) lettuce is better in MAP than in air. Green tissue had the lowest overall quality after air storage. In addition, there were no significant overall quality differences for red tissue stored under either condition. Texture and aroma were excellent for green tissue held in MAP, while it was unacceptable in air. Both atmospheres maintained texture of red and white tissues. Aroma was excellent for white tissue in both air and MAP. In contrast, aroma was only acceptable for red tissue stored in air.

#### ***Romaine lettuce***

Fresh-cut romaine should be stored at 34 to 38 °F (1 to 3 °C) before and after processing to assure quality and reduce potential for freezing of product during handling, distribution and storage. MAP exerted a beneficial effect on sensory quality by preventing pink discoloration. Compared to air, 3% O<sub>2</sub> + 6, 10, or 14% CO<sub>2</sub> MAP delayed development of tissue discoloration and increased shelf-life by about 50%. Of the three CO<sub>2</sub> levels, 10% was slightly more effective than 6 and 14% (Hamsa et al., 1996; Lopez-Galvez et al., 1996; Segall and Scanlon, 1996).

The aerobic plate counts of fresh-cut Romaine lettuce in MAP were decreased by 0.15 and 0.35 kGy gamma irradiation (Prakash et al., 2000). The 0.35 kGy treatment decreased aerobic plate counts by 1.5 logs and yeast/mold counts by 1 log. A combination of MAP and 0.35 kGy resulted in a 2 log reduction in microbial counts. These differences were maintained through 22 days of storage. Irradiation at 0.15 kGy caused smaller reductions in microbial counts. A 10% loss in firmness was observed at 0.35 kGy, while other sensory attributes such as color, generation of off-flavor, and appearance of visual defects were not affected. The control sample reached microbial counts of 10<sup>7</sup> in 14 days compared to the MAP samples that reached the same level in 17 days, and the combination samples (0.35 kGy and MAP) that took 28 days. Color, flavor, and odor were unaffected by irradiation, although an off-odor was obvious in MAP samples

after 21 days (17% CO<sub>2</sub> + < 1% O<sub>2</sub>). Low-dose irradiation increased shelf-life of cut Romaine lettuce by 2 to 5 days (Guner et al., 1997)

### **MUSHROOMS (sliced)**

Fresh-cut mushroom slices are white with no darkening and wrinkling. Mushroom slices should be stored at 34 to 38 °F (1 to 3 °C). Wrinkling and brown surface patches are due to excessive moisture loss (Roy et al., 1996). A CA of 10% CO<sub>2</sub> + 3% O<sub>2</sub> maintains quality, while 5% O<sub>2</sub> + 5% CO<sub>2</sub> at 5 °C slightly reduces respiration. More than 20% CO<sub>2</sub> accelerates tissue darkening, while < 3% O<sub>2</sub> may result in potential *botulinum* toxin production. Inoculation studies have demonstrated that *C. botulinum* type A will sporulate and produce endotoxin in 0.9 to 2% O<sub>2</sub> after 6 days at 75 to 79 °F (24 to 26 °C) (Sugiyama and Yang, 1975; Sugiyama and Rutledge, 1978). It is not recommended to use low O<sub>2</sub> MAP due to the inherent food safety risk.

#### **Respiration Rates: sliced**

Temperature	mg CO <sub>2</sub> kg <sup>-1</sup> h <sup>-1</sup>
0 °C	20 to 60
5 °C	39 to 88
10 °C	86 to 124

To get mL kg<sup>-1</sup> h<sup>-1</sup>, divide the mg kg<sup>-1</sup> h<sup>-1</sup> rate by 2.0 at 0 °C (32 °F), 1.9 at 10 °C (50 °F), and 1.8 at 20 °C (68 °F). To calculate heat production, multiply mg kg<sup>-1</sup> h<sup>-1</sup> by 220 to get BTU per ton per day or by 61 to get kcal per metric ton per day. Data from Gorny (1997).

### **ONIONS (Diced, slivered, rings, chunks)**

Fresh-cut onions should have no discoloration, skin and core. Bulbs should be dry, free of decay, firm and 3 to 4" (7.5 to 10 cm) in diameter. The core of whole onions should be 35 °F (1.5 °C) at receiving, and at 34 to 38 °F (1 to 3 °C) before and after processing to assure quality and reduce potential for freezing of product during handling, distribution and storage. The seed stems and new internal growth should be less than 10%; double centers and translucent scale should be less than 5%.

Whole onions are peeled and trimmed by machine or hand. Washing with chlorinated water can be done before or after processing. Bulbs for onion rings are washed with cold water at 32 °F (0 °C) before processing. However, diced and slivered onions are washed using chlorinated water after processing.

Browning, yellowing and development of translucence are major factors affecting the visual quality of diced onion (Blanchard et al., 1996). Cutting causes important biochemical changes in tissues, including development of sulfur aromatic volatiles. These compounds are produced enzymatically by hydrolysis of alliin (odorless derivatives of amino acids) by alliinase (Schwimmer and Weston, 1961; MacLoed, 1970).

Among the volatiles present in the aromatic profile of onion are oxides of disulfide, thiosulfates and propene disulfide, known bacteriostatics (Davidson et al, 1983). Their action on microorganisms may be due to the inhibition of respiratory enzymes containing thiol groups (Augusti, 1990).

Thiopropanal-S-oxide, a lacrimogenous factor, has antifungal properties (Sharmon et la., 1979) and may be involved in development of bitterness and browning of onion puree (Howard et al., 1994).

CA of 2 to 5% O<sub>2</sub> + 10 to 15% CO<sub>2</sub> decreases respiration and microbial proliferation, and retains sucrose and pungency of cut onions. Lowering the O<sub>2</sub> level can be beneficial for visual quality, but has no effect on aroma. Enrichment in CO<sub>2</sub> improves both qualities. Rapid reduction of O<sub>2</sub> in the package reduces discoloration. Browning can be slowed slightly by lowering O<sub>2</sub>, and more so by CO<sub>2</sub> enrichment. Langerak (1975) reported that freshly cut onion became brown and unacceptable after cooking when wrapped in perforated film, in comparison with onion wrapped in film that achieved MAP conditions more rapidly. Atmospheres containing 1 to 2% O<sub>2</sub> prevent the loss of pyruvic acid, an indicator of onion flavor intensity,

and reduce the rate of respiration of the fresh-cut tissue (Mencarelli et al., 1990).

The inclusion of potassium permanganate as an ethylene scrubber and sulfur volatiles in packages of diced onions improves sensory quality. A storage-life of 10 days is possible for fresh-cut onions packaged in polymeric bags without a gas flush and stored at 36 °F (2.2 °C). Addition of absorbents, potassium permanganate and clay, in packages increased shelf-life up to 18 days at 34 °F (1 °C) (Howard et al., 1994; Toivonen, 1997).

Yeasts and molds were virtually absent on diced onion. When present, their count did not increase beyond 10<sup>2</sup> CFU/g (Garg et al., 1990; Blanchard et al., 1996). Psychrotrophic and mesophilic flora increased throughout storage under all treatments. Their growth in diced onion was slowed under CA as compared to air (Blanchard et al., 1996).

Respiration rates vary significantly depending on how long whole onions were in cold storage prior to processing. CA (5% O<sub>2</sub> + 5% CO<sub>2</sub>) at 41 °F (5 °C) reduces respiration rate by 25%.

### Respiration Rates:

Temperature	Sliced (2 mm thick rings)	Diced (1 x 1 cm)
	(mg CO <sub>2</sub> kg <sup>-1</sup> h <sup>-1</sup> )	
2 °C	14.0	12.0
5 °C	23.4	15.6
10 °C	38.0	22.8
23 °C	126 to 131	90 to 99

To get mL kg<sup>-1</sup> h<sup>-1</sup>, divide the mg kg<sup>-1</sup> h<sup>-1</sup> rate by 2.0 at 0 °C (32 °F), 1.9 at 10 °C (50 °F), and 1.8 at 20 °C (68 °F). To calculate heat production, multiply mg kg<sup>-1</sup> h<sup>-1</sup> by 220 to get BTU per ton per day or by 61 to get kcal per metric ton per day. Data from Gorny 1997.

### PEPPER, Bell (Diced, Sliced)

Fresh-cut bell pepper should have no discoloration (darkening) or water soaking. Raw material should be received at < 45 °F (7 °C) and stored at 45 to 50 °F (7 to 10 °C), but stored at 34 to 40 °F (1 and 4 °C) after processing to assure quality and reduce potential for freezing of product during handling, distribution and storage. Defects that reduce overall visual quality includes darkening of the green or red pulp, brown discoloration of the cut surfaces, and decay. Whole peppers should be spray-washed before processing and washed after being cored and cut to reduce microbial population. Alternatively, cut pepper can be washed using a combination of open flume and close flume systems.

Although intact peppers are chilling sensitive, it is necessary to store fresh-cut red and green pepper at 32 to 41 °F (0 to 5 °C) to maintain visual quality (El-Bassuoni and Cantwell, 1994). Visual quality was maintained and few compositional changes occurred during 15 days of storage at 34 °F (1 °C) (Abe et al. 1991; Zhou et al., 1992). Full ripe or red peppers are less chilling sensitive than mature-green fruits of the same cultivar.

The recommended CA is 3% O<sub>2</sub> + 5 to 10% CO<sub>2</sub>. Levels of CO<sub>2</sub> > 10% can result in tissue darkening and softening. Shelf-life and sensitivity to CO<sub>2</sub> injury varies significantly among cultivars. Air + 10% CO<sub>2</sub> maintains quality at both 41 and 50 °F (5 and 10 °C). Aroma and texture decrease after 6 days in 10% CO<sub>2</sub>, but were maintained in air at 0°C. Storage in high CO<sub>2</sub> at 41 °F (5 °C) and air at 32 °F (0 °C) retard decay development. Atmospheres with 5% CO<sub>2</sub> help maintain quality, but are not as effective as 10% CO<sub>2</sub>. Fresh-cut pepper pieces that were ripening or were already fully red ripe had better shelf-life than pieces from mature-green fruit (Lopez-Galvez et al., 1997). They also found that visual quality was best maintained in air or CA at 32 °F (0 °C), with important quality loss observed by day 9 at 41 °F (5 °C) and by day 3 at 50 °F (10 °C). Decay is caused by both fungi (*Alternaria* and *Botrytis* spp.) and soft-rot causing bacteria. Diced tissue was more susceptible to decay than slices (Lopez-Galvez et al., 1997).

**Respiration Rates:**

Temperature	Sliced (5 cm)	Diced (1 x 1 cm)
	(mg CO <sub>2</sub> kg <sup>-1</sup> h <sup>-1</sup> )	
0 °C	2 to 6	2 to 9
5 °C	4 to 7	6 to 12
10 °C	6 to 13	11 to 18

To get mL kg<sup>-1</sup> h<sup>-1</sup>, divide the mg kg<sup>-1</sup> h<sup>-1</sup> rate by 2.0 at 0 °C (32 °F), 1.9 at 10 °C (50 °F), and 1.8 at 20 °C (68 °F). To calculate heat production, multiply mg kg<sup>-1</sup> h<sup>-1</sup> by 220 to get BTU per ton per day or by 61 to get kcal per metric ton per day. Data from Gorny 1997.

**POTATO (sticks, diced, sliced, peeled)**

Fresh-cut potatoes should be firm and without brown discoloration. Enzymatic browning is a major problem in the discoloration of peeled or fresh-cut potatoes. Hand-peeling and lye-peeling result in good quality, while abrasion-peeling is undesirable for fresh potatoes. Anti-browning dips like 0.5% L-cysteine + 2% citrate, in combination with MAP, are needed to prevent browning of whole peeled potatoes.

Pre-treatment with heated ascorbate and citrate solutions, prior to a browning inhibitor dip containing 4% ascorbate, 1% citrate and 1% sodium pyrophosphate, has greatly extended storage-life over that after the dip alone (Sapers and Miller, 1995; Sapers et al., 1997). However, under some conditions, such pre-treatment induced changes in the cooked product that result in surface firming (case hardening) and partial separation of the case-hardened superficial layer from underlying tissue during slicing. Furthermore, case-hardened potatoes form large lumps during mashing (Martin et al., 1999). The recommended storage temperature is 32 °F (0 °C). Shelf-life of minimally-processed products can be extended to 3 weeks under CA and refrigeration.

Dipping potato strips in chlorine solutions (100 and 300 ppm) result in higher microbial populations during storage, while potatoes treated with anti-browning solutions combined with MAP showed only a slight increase. MAP had no significant effect on microbial populations compared to non-packaged samples. The predominant organisms were *Pseudomonas fluorescens* (Gunes et al., 1997).

The recommended CA is 1 to 3% O<sub>2</sub> + 6 to 9% CO<sub>2</sub>. Reducing O<sub>2</sub> level to 10 or 3% + 10% CO<sub>2</sub> at 2 °C reduces respiration of potato sticks by 50 to 75%. However, 5% O<sub>2</sub> + 5% CO<sub>2</sub> at 5°C increases respiration of sticks during storage.

Active modification of package atmospheres is necessary to prevent browning and achieve extended shelf-life. MAP alone will not prevent browning (Gurbuz and Lee, 1997). Vacuum packaging of fresh-cut potatoes may create anaerobic conditions that allow growth of *Clostridium botulinum*.

**Respiration Rates:**

Temperature	Whole Peeled	Halves	Slices (2 mm)	Sticks (0.95 x 0.95 cm)
	(mg CO <sub>2</sub> kg <sup>-1</sup> h <sup>-1</sup> )			
2 °C	6.0 to 8.0	8.0	10.0 to 12.0	12.2
5 °C	7.8	7.8 to 9.8	11.7 to 15.6	-
10 °C	17.1 to 19.0	21 to 22.8	38.0	-
23 °C	54 to 63	81	-	117 to 126

To get mL kg<sup>-1</sup> h<sup>-1</sup>, divide the mg kg<sup>-1</sup> h<sup>-1</sup> rate by 2.0 at 0 °C (32 °F), 1.9 at 10 °C (50 °F), and 1.8 at 20 °C (68 °F). To calculate heat production, multiply mg kg<sup>-1</sup> h<sup>-1</sup> by 220 to get BTU per ton per day or by 61 to get kcal per metric ton per day. Data from Gorny (1997).

**RUTABAGA (cubed, shredded, peeled)**

Fresh-cut rutabaga should have uniform yellow flesh without decay, bruises or discoloration. Raw material

should arrive at < 45 °F (7 °C), while raw and fresh-cut product should be stored at 34 to 38 °F (1 to 3 °C) before and after processing to assure quality and reduce potential for freezing during handling, distribution and storage. A CA of 5% CO<sub>2</sub> + 5% O<sub>2</sub> is slightly beneficial and reduces respiration at 41 °F (5 °C).

**Respiration Rates:**

Temperature	1/4 Whole, Peeled	Shredded (2 mm) (mg CO <sub>2</sub> kg <sup>-1</sup> h <sup>-1</sup> )	Diced (1 x 1 cm)
2 °C	8	16	18
5 °C	12	20	25
10 °C	15	42	42 to 48
23 °C	45	72 to 81	153 to 162

To get mL kg<sup>-1</sup> h<sup>-1</sup>, divide the mg kg<sup>-1</sup> h<sup>-1</sup> rate by 2.0 at 0 °C (32 °F), 1.9 at 10 °C (50 °F), and 1.8 at 20 °C (68 °F). To calculate heat production, multiply mg kg<sup>-1</sup> h<sup>-1</sup> by 220 to get BTU per ton per day or by 61 to get kcal per metric ton per day. Data from Gorny (1997).

**SPINACH (whole leaves, cut leaves)**

Fresh-cut spinach should be green without any decay or bruise. Raw material should arrive at 35 °F (1.5 °C). Fresh and processed products should be stored at 34 to 38 °F (1 to 3 °C) before and after processing to assure quality and reduce potential for freezing of product during handling, distribution and storage.

Storage in 0.8 to 3% O<sub>2</sub> + 8 to 10% CO<sub>2</sub> is beneficial. Spinach stored in 0.8% O<sub>2</sub> at 20°C (68 °F) had a lower respiration rate, superior appearance, and better taste than spinach leaves stored in air (Izumi et al., 1997). Atmospheres of 0.8% O<sub>2</sub> or 0.8% O<sub>2</sub> + 10% CO<sub>2</sub> reduced the number of aerobic mesophilic and psychrotrophic microorganisms on fresh-cut spinach leaves by 10- to 100-fold, compared to air-stored leaves at 41 °F (5 °C); however, there was no effect at 50 °F (10 °C). These atmospheres had no effect on texture changes during storage at 41 °F (5 °C) for 9 days or at 50 °F (10 °C) for 7 days (Babic and Watada, 1996).

An atmosphere of 4% O<sub>2</sub> + 9% CO<sub>2</sub> reduced the loss of ascorbic acid by 50% compared to air (Burgheimer et al., 1967). Increasing CO<sub>2</sub> concentrations up to 13% caused a rapid decrease in ascorbic acid and development of off-odors that were not acceptable after 1 week storage at 45 °F (7 °C) (McGill et al., 1966). Chlorophyll loss can be reduced by 50% with ethylene scrubbing. Low O<sub>2</sub> (0.8%) results in a slightly reduced respiration rate of cleaned spinach leaves.

Microbiological and sensory changes occur in all packaged spinach samples, being more pronounced at 50 °F (10 °C) than at 40 °F (4 °C). Spinach leaves harbor high numbers of mesophilic, psychrotrophic and Pseudomonadaceae bacteria (Babic and Watada, 1996). The pectinolytic species *Pseudomonas fluorescens* is probably the major spoilage agent of fresh-cut spinach. Microbial populations on fresh-cut spinach leaves increase during storage in air and CA at 41 and 50 °F (5 and 10 °C). Populations of Enterobacteriaceae and micrococcaceae are not greatly affected by storage atmosphere.

**Respiration Rate: whole leaves**

Temperature	mg CO <sub>2</sub> kg <sup>-1</sup> h <sup>-1</sup>
0 °C	6.0 to 14.0
5 °C	11.7 to 23.4

To get mL kg<sup>-1</sup> h<sup>-1</sup>, divide the mg kg<sup>-1</sup> h<sup>-1</sup> rate by 2.0 at 0 °C (32 °F), 1.9 at 10 °C (50 °F), and 1.8 at 20 °C (68 °F). To calculate heat production, multiply mg kg<sup>-1</sup> h<sup>-1</sup> by 220 to get BTU per ton per day or by 61 to get kcal per metric ton per day. Data from Gorny (1997).

### **SWEET POTATO (shredded, sliced)**

Fresh-cut sweet potatoes should be firm and without discoloration. Whole sweet potatoes are washed with water before peeling and shredding. Cut sweet potatoes are then washed with 100 ppm sodium hypochlorite, rinsed, and centrifuged to remove excess water. Product quality remains acceptable during 2 weeks storage at 32 to 40 °F (0 to 4 °C) in polymeric bags. Color of shredded sweet potato darkens and firmness declines during storage.  $\beta$ -Carotene remains stable at 7.8 mg per 100 g at 40 °F (4 °C). Vitamin C remains stable at 7 mg per 100 g for 1 week and then decreases. A 6.5% CO<sub>2</sub> + 12% O<sub>2</sub> atmosphere is slightly beneficial.

### **TOMATO (Sliced, Diced)**

Fresh-cut tomatoes should have consistent red color and firm texture. Sliced tomatoes have no gel loss from the seed cavities. Diced tomatoes have no seeds and stem pieces. 'Roma' tomatoes are often used for diced tomatoes; the ripening stage should be 5 or higher (USDA standard). The ripening stage of round tomatoes for slicing should be 5.5 or higher. Fruit should be firm and have small seed cavities. For diced tomato products, 'Roma' tomatoes should be spray-washed with chlorinated water before cutting. After cutting, an open fluming system with 0.25% CaCl<sub>2</sub> and 100 ppm chlorine is used to wash and remove seeds. For sliced tomato products, whole tomatoes are dipped in 200 ppm chlorinated water for more than 1 min before slicing. Various automated slicing machines have been developed for tomatoes. Sliced tomatoes can be packed with stacking, shingling or soldier style.

Although tomatoes are chilling sensitive, they can be stored at 32 to 41 °F (0 to 5 °C) for a few days before processing to retard softening. A CA of 3% O<sub>2</sub> + 3% CO<sub>2</sub> will delay ripening, as well as SSC and TA losses (Mencarelli and Saltveit, 1988). Water soaking that produces translucent tissue, textural changes, and softening reduces the quality of diced tomatoes during storage. In addition to these deleterious changes, quality of tomato slices can be reduced by seed germination and loss of locular gel. The high acidity of tomato products suppresses microbial growth, but growth of yeasts and molds can reduce quality in storage.

#### **Respiration Rate:**

Temperature	mg CO <sub>2</sub> kg <sup>-1</sup> h <sup>-1</sup>
20 °C	57.6 to 93.6

To get mL kg<sup>-1</sup> h<sup>-1</sup>, divide the mg kg<sup>-1</sup> h<sup>-1</sup> rate by 2.0 at 0 °C (32 °F), 1.9 at 10 °C (50 °F), and 1.8 at 20 °C (68 °F). To calculate heat production, multiply mg kg<sup>-1</sup> h<sup>-1</sup> by 220 to get BTU per ton per day or by 61 to get kcal per metric ton per day.

### **ZUCCHINI (Slices)**

Fresh-cut zucchini should have a dark green peel, white tissue and a crisp texture. Raw material should be received at < 55 °F (13 °C) and stored at 41 to 50 °F (5 to 10 °C). Storage should be at 32 to 41 °F (0 to 5 °C) after processing. A CA of 0.25 to 1% O<sub>2</sub> is beneficial. Lowering O<sub>2</sub> to < 0.5% at 41 °F (5 °C) reduces respiration by 50%; respiration is reduced by 80% at 50 °F (10 °C). Sliced zucchini develops water soaked areas (chilling injury) at 32 °F (0 °C) and brown discoloration at 41 to 50 °F (5 to 10 °C) which increases with storage duration. Zucchini slices can be dipped in solutions of CaCl<sub>2</sub> alone or with NaOCl. Calcium treatments reduce development of decay and total microbial growth, and ascorbate loss.

#### **Respiration Rates:**

Temperature	mg CO <sub>2</sub> kg <sup>-1</sup> h <sup>-1</sup>
5 °C	21.1
10 °C	19.4 to 49.4

To get  $\text{mL kg}^{-1} \text{h}^{-1}$ , divide the  $\text{mg kg}^{-1} \text{h}^{-1}$  rate by 2.0 at  $0\text{ }^{\circ}\text{C}$  ( $32\text{ }^{\circ}\text{F}$ ), 1.9 at  $10\text{ }^{\circ}\text{C}$  ( $50\text{ }^{\circ}\text{F}$ ), and 1.8 at  $20\text{ }^{\circ}\text{C}$  ( $68\text{ }^{\circ}\text{F}$ ). To calculate heat production, multiply  $\text{mg kg}^{-1} \text{h}^{-1}$  by 220 to get BTU per ton per day or by 61 to get kcal per metric ton per day. Data from Gorny (1997).

### References:

- Abe, K., K. Yoshimura, Y.-F. Zhou, and T. Iwata. 1991. Studies on physiological and chemical changes of partially processed sweet pepper fruit. II. Changes of chemical compounds in cut surface of sweet pepper in relation to difference of storability by shredding direction. *J. Jap. Soc. Cold Preserv. Food* 17:146-151.
- Augusti, K.T. 1990. Therapeutic and medicinal values of onions and garlic. In: J.L. Brewster and H.D. Rabinowitch (eds) *Onions and allied crops. Vol. III: Biochemistry. Food Science and Minor Crops.* CRC Press, Boca Raton FL, pp. 93-108.
- Avena-Bustillos, R.D.J. J.M. Krochta, and M.E. Saltveit. 1997. Water vapor resistance of red delicious apples and celery sticks coated with edible caseinate-acetylated monoglyceride films. *J. Food Sci.* 62:351-354.
- Avena-Bustillos, R.J., L.A. Cisneros-Zevallos, J.M. Krochta, and M.E. Saltveit. 1994. Application of casein-lipid edible film emulsions to reduce white blush on minimally processed carrots. *Postharv. Biol. Technol.* 4:319-329.
- Babic, I. and A.E. Watada. 1996. Microbial populations of fresh-cut spinach leaves affected by CAs. *Postharv. Biol. Technol.* 9:187-193.
- Beaulieu, J.C., F.A.R. Oliveira, T.D. Fernandes, S.C. Fonseca and J.K. Brecht. 1997. Fresh-cut kale: quality assessment of Portuguese store-supplied product for development of a MAP system. *Proc. Vol. 5: Fresh-Cut Fruits and Veg. and MAP.* J. Gorny (ed) Davis CA, pp. 145-151.
- Beuchat, L.R. and R.E. Brackett. 1990. Survival and growth of *Listeria monocytogenes* on lettuce as influenced by shredding, chlorine treatment, modified atmosphere packaging and temperature. *J. Food. Sci.* 55: 755-758.
- Blanchard, M., F. Caistaingne, C. Willemot and J. Makhlof. 1996. Modified atmosphere preservation of freshly prepared diced yellow onions. *Postharv. Biol. Technol.* 9:173-185.
- Bolin, H.R. and C.C. Huxsoll. 1991. Control of minimally processed carrot (*Daucus carota*) surface discoloration caused by abrasion peeling. *J. Food Sci.* 56:416-418.
- Brecht, J.K. 1995. Physiology of lightly processed fruits and vegetables. *HortScience* 30: 18-21.
- Burgheiner, F., J.N. Macgill, A.I. Nelson and M.P. Steinberg. 1967. Chemical changes in spinach stored in air and CA. *Food Technol.* 21:109-111.
- Burlo-Carbonell, F.M., A.A. Carbonell-Barrachina, D. Valero, D. Martinez-Romero and F. Martinez-Sanchez. 2000. Acid treatment and edible coatings can improve preservation in fresh garlic. *Tech. Prog. Book Abstr., 2000 IFT Annual Meeting.* pp. 133.
- Cabezas, A. and D.G. Richardson. 1997. Modified atmosphere packaging of broccoli florets of temperature and package type. *Vol. 5: Fresh-Cut Fruits and Veg. and MAP.* J. Gorny (ed) Davis CA, pp. 8-15.
- Carlin, F., C. Nguyen-The, Y. Cudennec and M. Reich. 1989. Microbial spoilage of fresh, "ready-to-use" grated carrots. *Sciences des Aliments.* 9:371-386.
- Carlin, F., C. Nguyen-The, G. Hilbert, and Y. Chambroy. 1990. Modified atmosphere packaging of fresh, "ready-to-use" grated carrots in polymeric films. *J. Food. Sci.* 55:1033-1038.
- Castaner, M., M.I. Gil, F. Ferreres, F. Artes, and F.A. Tomas-Barberan. 1997. Effect of modified atmospheres on anthocyanin stability and other phenolic metabolites during storage of minimally processed red lettuce. *Vol. 5, Fresh-cut Fruits and Vegetables and MAP.* J. Gorny (ed) Davis CA, pp. 134-138.
- Cisneros-Zevallos, L., M.E. Saltveit and J.M. Krochta. 1995. Mechanism of surface white discoloration of peeled (minimally processed) carrots during storage. *J. Food Sci.* 60:320-323, 333.
- Cisneros-Zevallos, L.A., M.E. Saltveit and J.M. Krochta. 1997. Hygroscopic coatings control surface white

- discoloration of peeled (minimally processed) carrots during storage. *J. Food Sci.* 62:363-366.
- Davidson, P.M., L.S. Post, A.L. Branen and A.R. McCordy. 1983. Naturally occurring and miscellaneous food antimicrobials. In: A.L. Bradeo and P.M. Davidson (eds) *Antimicrobials in Foods*, Marcel Dekker, NY, pp. 275-276.
- El-Bassuoni, R. and M. Cantwell. 1994. Low temperatures and CAs maintain quality of fresh cut bell pepper. *HortScience* 29:448.
- Galetti, L., H. Berger, V. Escalona, C. Saenz, and E. Araya. 1997. Modified atmospheres and ascorbic acid for minimum process of carrot slices. *Proc. Vol. 5: Fresh-Cut Fruits and Veg. and MAP. J. Gorny (ed) Davis CA*, pp. 139-144.
- Garg, N., J.I. Churey and D.F. Splinsioester. 1990. Effect of processing conditions on the microflora of fresh-cut vegetables *J. Food Prot.* 53:701-703.
- Gariepy, Y., G.S.V. Raghavan and R. Theriault. 1986. CA storage of celery with the silicone membrane system. *Int. J. Refrig.* 9:234-239.
- Gil-Maria, I., M. Castaner, F. Ferreres, F. Artes, and B.F.A. Tomas. 1998. Modified-atmosphere packaging of minimally processed "Lollo Rosso" (*Lactuca sativa*): Phenolic metabolites and quality changes. *Zeitschrift Lebensmittel Untersuchung Forschung.* 206:350-354.
- Gorny, J.R. 1997. A summary of CA and Ma requirements and recommendations for fresh-cut (minimally processed) fruits and vegetables. Vol 5, *Fresh-cut Fruits and Vegetables and MAP. J. Gorny (ed) Davis CA*, pp. 30-67.
- Gunes, G., D.F. Splittstoesser, and C.Y. Lee. 1997. Microbial quality of fresh potatoes: effect of minimal processing. *J. Food Prot.* 60:863-866.
- Gurbuz, G. and C.Y. Lee. 1997. Color of minimally processed potatoes as affected by modified atmosphere packaging and anti-browning agents. *J. Food Sci.* 62:572-575.
- Hagenmaier, R.D. and R.A. Robert. 1998. Microbial population of shredded carrot in modified atmosphere packaging as related to irradiation treatment. *J. Food Sci.* 63:162-164.
- Hamza, F., F. Castaigne, C. Willemot, G. Doyon and J. Makhlof. 1996. Storage of minimally processed romaine lettuce under CA. *J. Food Qual.* 19:177-188.
- Hiroaki, K., U. Masayuki and O. Yutaka. 1993. Storage of shredded cabbage under dynamically CA of high O<sub>2</sub> and high CO<sub>2</sub>. *Biosci. Biotech. Biochem.* 57:1049-1052.
- Howard, L.R. and T. Dewi. 1995. Sensory, microbiological and chemical quality of mini-peeled carrots as affected by edible coating treatment. *J. Food Sci.* 60:142-144.
- Howard, L.R. and T. Dewi. 1996. Minimal processing and edible coating effects on composition and sensory quality of mini-peeled carrots. *J. Food Sci.* 61:643-646.
- Howard, L.R., K.S. Yoo, L.M. Park, and G.H. Miller Jr. 1994. Quality changes in diced onions stored in film packages. *J. Food Sci.* 59:110-112.
- Izumi, H., T. Nonaka and T. Muraoka. 1997. Physiology and quality of fresh-cut spinach stored in low O<sub>2</sub> CAs at various temperature. Vol. 5, *Fresh-cut Fruits and Vegetables and MAP. J. Gorny (ed) Davis CA*, pp. 130-133.
- Izumi H., A.E. Watada, and W. Douglas. 1996a. Optimum O<sub>2</sub> or CO<sub>2</sub> atmosphere for storing broccoli florets at various temperatures. *J. Amer. Soc. Hort. Sci.* 121:127-131.
- Izumi, H., A.E. Watada, N.P. Ko, and W. Douglas. 1996b. CA storage of carrot slices, sticks and shreds. *Postharvest Biol. Technol.* 9:165-172.
- Johnson, C.E. and J.H. von Elbe. 1974. Extension of postharvest storage-life of sliced celery. *J. Food Sci.* 39:678-680.
- Keteleer, A. C. De Mulder, A. Hermans and P. Tobback. 1993. Biplot presentation of the physiological-biochemical characterization and the sensorial quality of minimally processed leek stored under modified atmosphere. *Acta Hort.* 343:345-346.
- Krahn, T.R. 1977. Improving to keeping quality of cut head lettuce. *Acta Hort.* 62:79-92.
- Langerak, D.I. 1975. The influence of irradiation and packaging on the keeping quality of prepared cut endive, chicory and onions. *Acta Alim.* 4:123-138.

- Li, P. and M.M. Barth. 1998. Impact of edible coatings on nutritional and physiological changes in lightly-processed carrots. *Postharvest. Biol. Technol.* 14:51-60.
- Loaiza-Velarde, J.G., F.A. Tomás-Barberá and M.E. Saltveit. 1997. Effect of intensity and duration of heat-shock treatments on wound-induced phenolic metabolism in Iceberg lettuce. *J. Amer. Soc. Hort. Sci.* 122:873-877.
- Loaiza-Velarde, J.G. and M.E. Saltveit. 2001. Heat shocks applied either before or after wounding reduce browning of lettuce leaf tissue. *J. Amer. Soc. Hort. Sci.* 126:227-234.
- Loaiza-Velarde, J.G., M.E. Mangrich, R. Campos-Vargas and M.E. Saltveit. 2002. Heat-shock reduces browning of fresh-cut celery petioles. *Postharv. Biol. Technol.* (In Press)
- Lopez-Galvez, G., R. El-Bassuoni, X. Nie, and N. Cantwell. 1997. Quality of red and green fresh-cut peppers stored in CAs. Vol. 5, *Fresh-Cut Fruits and Veg. and MAP.* J. Grony (ed) Davis CA, pp. 152-157.
- Lopez-Galvez, G., M. Saltveit and M. Cantwell. 1996. The visual quality of minimally processed lettuces stored in air or CA with emphasis on romaine and iceberg types. *Postharv. Biol. Technol.* 8:179-190.
- MacLeod, A.L. 1970. The chemistry of vegetable flavor. *Flav. Ind.*, October, pp. 665-672.
- Makhlouf, J., F. Castaign, J. Arul, C. Willemot, and A. Gosselin. 1989. Long-term storage of broccoli under CA. *HortScience* 24:637-639.
- Martin, S.T., G.M. Sampers and R.L. Miller. 1999. Process for inhibiting enzymatic browning and maintaining textural quality of fresh peeled potatoes. U.S. Patent No. 5912034.
- Mateos, M., D. Ke, M. Cantwell and A.A. Kader. 1993. Phenolic metabolism and ethanolic fermentation of intact and cut lettuce exposed to CO<sub>2</sub>-enriched atmospheres. *Postharv. Biol. Technol.* 3:225-233.
- McDonald, R.E., L.A. Risse, and C.R. Barmore. 1990. Bagging chopped lettuce in permeability films. *HortScience* 25:671-673.
- McGill, J.N., A.I. Nelson and M.P. Steinberg. 1966. Effects of modified storage atmospheres on ascorbic acid and other quality characteristics of spinach. *J. Food Sci.* 31:510-517.
- Mencarelli, F., D. De Santic, R. Massantini and M. Contini. 1990. Low O<sub>2</sub> concentrations use for the short-term onion slice storage. *Industrie Alimentari.* 29:982-984.
- Mencarelli, F. and M.E. Saltveit Jr. 1988. Ripening of mature-green tomato fruit slices. *J. Amer. Soc. Hort. Sci.* 113:742-745.
- Nguyen-the, C. and J. Prunier. 1989. Involvement of *Pseudomonas* in 'ready-to-use' salad deterioration, *Int. J. Food Sci. Technol.* 24:47-56.
- Omary, M., R. Testin, S. Barefoot and J. Rushing. 1993. Packaging effects on growth of *Listeria innocua* in shredded cabbage. *J. Food Sci.* 58:623-626.
- Peiser, G., G. Lopez-Galvez and M. Cantwell. 1997. Changes in aromatic volatiles during storage of packed salad products. *Proc. 7<sup>th</sup> Intl. CA Research Conference, July 1997, Davis CA.*
- Pirovani, M.E., D.R. Guenes, A.M. Piagentini and J.H.D. Pentima. 1997. Storage quality of minimally processed cabbage packaged in plastic films. *J. Food Qual.* 20:381-389.
- Poulsen, N., A.S. Johnson and J.N. Sorensen. 1995. Influence of growth conditions on the value of crisphead lettuce, 4: Quality changes during storage. *Plant Foods Hum. Nutrl* 47:157-162.
- Poulsen, N., J.N. Sorensen and A.S. Johnson. 1994. Influence of growth conditions on the value of crisphead lettuce, 2: weight losses during storage as affected by nitrogen, plant age and cooling system. *Plant Foods Hum. Nutrl* 46:13-18.
- Prakash, A. A.R. Guner, F. Caporaso, and D.M. Foley. 2000. Effects of low-dose Gamma irradiation on the shelf-life and quality characteristics of cut romaine lettuce packaged under modified atmosphere. *J. Food Sci.* 65:549-553.
- Robbs, P.G., J.A. Bartz, G. McFie, and N.C. Hodge. 1996. Causes of decay of fresh-cut celery. *J. Food Sci.* 61:444-448.
- Roy, S., R.C. Anantheswaran and R.B. Beelman. 1996. Modified atmosphere and modified humidity packaging of fresh mushrooms. *J. Food Sci.* 61:391-397.
- Saltveit, M.E. 1997. Physical and physiological changes in minimally processed fruits and vegetables. In:

- Phytochemistry of fruit and vegetables. F.A. Tomás-Barberán and R.J. Robins (eds), pp. 205-220. Proceedings of the phytochemical society of Europe 41. Oxford Univ. Press, NY.
- Saltveit M.E. 2000. Wound induced changes in phenolic metabolism and tissue browning are altered by heat shock. *Postharv. Biol. Technol.* 21:61-69.
- Saltveit, M.E. and M.E. Mangrich. 1996. Use of density measurements to study the effect of excision, storage, abscisic acid and ethylene on the development of pithiness in celery (*Apium graveolens* L.) petioles. *J. Amer. Soc. Hort. Sci.* 121:137-141.
- Sapers, G.M., P.H. Cooke, A.E. Heidel, S.T. Martin and R.L. Miller. 1997. Structural changes related to texture of pre-peeled potatoes. *J. Food Sci.* 62:797-803.
- Sapers, G.M. and R.L. Miller. 1995. Heated ascorbic citric acid solution as browning inhibitor treatment for pre-peeled potatoes. *J. Food Sci.* 60:762-766.
- Schwimmer, S. and W.J. Weston. 1961. Enzymatic development of pyruvic in onion as a measure of pungency. *Ag. Food Chem.* 9:301-304.
- Segall, K.I. and M.G. Scanlon. 1996. Design and analysis of a modified-atmosphere package for minimally processed romaine lettuce. *J. Amer. Soc. Hort. Sci.* 121:722-729.
- Sharmon, A., G.M. Tewari, A.J. Shrikhande, S.R. Padwal-Desai and C. Brandyopadhyay. 1979. Inhibition of aflatoxin-producing fungi by onion extract. *J. Food Sci.* 44:1545-1547.
- Smyth-Anne, B., J. Song. And C. Cameron-Author. 1998. Modified atmosphere packaged cut iceberg lettuce: effect of temperature and O<sub>2</sub> partial pressure on respiration and quality. *J. Ag. Food. Chem.* 46:4556-4562.
- Sorensen, J.N., A.S. Johnnsen and N. Poulsen. 1994. Influence of growth conditions on the value of crisphead lettuce, 1: marketable and nutritional quality as affected by nitrogen supply, cultivar and plant age. *Plant Foods Human Nutr.* 46:1-12.
- Steward, J.K. and M. Uota. 1971. Carbon dioxide injury and market quality of lettuce held in CAs. *J. Amer. Soc. Hort. Sci.* 96:27-30.
- Stewart, J.K. and M.J. Ceponis. 1968. Effect of transit temperatures and modified atmospheres on market quality of lettuce shipped in nitrogen-refrigerated trailers. *USDA Ag. Mkt. Res. Rpt. No.* 832.
- Sugiyama, H. and K.S. Rutledge. 1978. Failure of *Clostridium botulinum* to grow in fresh mushrooms packaged in plastic film overwraps with holes. *J. Food Prot.* 41:348-350.
- Sugiyama, H. and K.H. Yang. 1975. Growth potential of *Clostridium botulinum* in fresh mushrooms packaged in semi-permeable plastic film. *App. Microbiol.* 30:964-969.
- Tatsumi, Y., A.E. Watada, W.P. Wergin. 1991. Scanning electron microscopy of carrot stick surface determines cause of white translucent appearance. *J. Food Sci.* 56:1357-1359.
- Toivonen, P.M.A. 1997. Quality changes in packaged, diced onions (*Alliumcepa* L.). Vol. 5, Fresh-cut Fruits and Veg. and MAP. J. Gorny (ed) Davis CA, pp. 1-6.
- Watada, A.E., N.P. Ko and D.A. Minott. 1996. Factors affecting quality of fresh-cut horticultural products. *Postharv. Biol. Technol.* 9:115-125.
- Zhou, Y.F., K. Abe, and T. Iwata. 1992. Effect of shredding modes on the deterioration of the quality of partially processed pepper fruits. *Nippon Shokuhin Kogyo Gakkaishi* 39:161-166.