

Chilling and Freezing Injury

Chien Yi Wang

Wang is formerly with the Produce Quality and Safety Laboratory, Henry A. Wallace Beltsville Agricultural Research Center, Agricultural Research Service, U.S. Department of Agriculture, Beltsville, MD. He is now retired.

Chilling Injury

Many fruits, vegetables, and ornamentals of tropical or subtropical origin are sensitive to low temperatures (Paull 1990). These crops are injured after a period of exposure to chilling temperatures below 10 to 15 °C (50 to 59 °F) but above their freezing points (Lyons 1973, Wang 1990). Certain horticultural crops of temperate origin are also susceptible to chilling injury (Bramlage and Meir 1990). Those temperate crops, in general, have lower threshold temperatures of <5 °C (41 °F). At these chilling temperatures, the tissues weaken because they are unable to carry on normal metabolic processes. Various physiological and biochemical alterations and cellular dysfunctions occur in chilling-sensitive species in response to chilling stress (Wang 1982, Wang and Adams 1982, Raison and Orr 1990). When chilling stress is prolonged, these alterations and dysfunctions will lead to the development of a variety of chilling injury symptoms such as surface lesions, internal discoloration, water-soaking of the tissue, and failure to ripen normally (Saltveit and Morris 1990). Often, products that are chilled will still look sound when remaining in low temperatures. However, symptoms of chilling injury become evident shortly after they are moved to warmer temperatures. Fruits and vegetables that have been chilled may be particularly susceptible to decay. Weak pathogens such as *Alternaria* spp., which do not grow readily on healthy tissues, can attack tissues that have been weakened by low-temperature exposure (McColloch and Worthington 1952, McColloch 1962).

Both temperature and duration of exposure are involved in the development of chilling injury. Damage may occur in a short time if temperatures are considerably below the threshold level, but a product may be able to withstand temperatures a few degrees into the critical zone for a longer time before injury becomes irreversible. Maturity at harvest and degree of ripeness are important factors in determining chilling sensitivity in some fruits like avocados (Kosiyachinda and Young 1976), honeydew melons (Lipton 1978), and tomatoes (McColloch et al. 1966). The effects of chilling are cumulative in some commodities. Low temperatures in transit, or even in the field shortly before harvest, add to the total effects of chilling that occur in cold storage.

Treatments shown to alleviate chilling injury include intermittent warming; high- or low-temperature preconditioning; CA storage; pretreatments with ethylene, abscisic acid, methyl jasmonate, and other natural compounds; calcium or other chemical applications; hypobaric storage; waxing; film packaging; and genetic manipulation (Ryall and Lipton 1979, Wang 1993, 1994, Meir et al. 1996).

Chilling injury is discussed more specifically under each commodity. Many of the commodities susceptible to chilling injury are listed in table 1 with threshold temperatures and some of the symptoms.

Table 1. Fresh produce susceptible to chilling injury when stored at low but nonfreezing temperatures

Commodity	Lowest Safe Temperature		Symptoms of injury when stored between 0 °C and safe temperature*
	° C	° F	
Apples—certain cultivars	2-3 [†]	36-38	Internal browning, brown core, soggy breakdown, soft scald
Asparagus	0-2	32-36	Dull, gray-green, limp tips
Atemoya	4	39	Skin darkening, failure to ripen, pulp discoloration
Avocados	4.5-13 [†]	40-55	Grayish-brown discoloration of flesh
Bael	3	38	Brown spots on skin
Bananas	11.5-13 [†]	53-56	Dull color when ripened
Bean (lima)	1-4.5	34-40	Rusty brown specks, spots or areas
Bean (snap)	7 [†]	45	Pitting and russeting
Breadfruit	7-12	45-53	Abnormal ripening, dull brown discoloration
Choyote	5-10	41-50	Dull brown discoloration, pitting, flesh darkening
Cranberries	2	36	Rubbery texture, red flesh
Cucumbers	7	45	Pitting, water-soaked spots, decay
Eggplants	7	45	Surface scald, alternaria rot, blackening of seeds
Ginger	7	45	Softening, tissue breakdown, decay
Guavas	4.5 [†]	40	Pulp injury, decay
Grapefruit	10 [†]	50	Scald, pitting, watery breakdown
Jicama	13-18	55-65	Surface decay, discoloration
Lemons	11-13 [†]	52-55	Pitting, membranous staining, red blotch
Limes	7-9	45-48	Pitting, turning tan with time
Lychee	3	38	Skin browning
Mangos	10-13 [†]	50-55	Grayish scald-like discoloration of skin, uneven ripening
Mangosteen	4-8	39-47	Hardening and browning of the cortex
Melons			
Cantaloupe	2-5 [†]	36-41	Pitting, surface decay
Honeydew	7-10	45-50	Reddish-tan discoloration, pitting, surface decay, failure to ripen
Casaba	7-10	45-50	Pitting, surface decay, failure to ripen
Crenshaw and Persian	7-10	45-50	Pitting, surface decay, failure to ripen
Okra	7	45	Discoloration, water-soaked areas, pitting, decay
Olive, fresh	7	45	Internal browning
Oranges	3 [†]	38	Pitting, brown stain

Papayas	7	45	Pitting, failure to ripen, off flavors, decay
Passion fruit	10	50	Dark red discoloration on skin, loss of flavor, decay
Peppers, sweet	7	45	Sheet pitting, alternaria rot on pods and calyxes, darkening of seeds
Pineapples	7-10 [†]	45-50	Dull green when ripe, internal browning
Pomegranates	4.5	40	Pitting, external and internal browning
Potatoes	3 [†]	38	Mahogany browning, sweetening
Pumpkins and hardshell squash	10	50	Decay, especially alternaria rot
Rambutan	10	50	Darkening of exocarp
Sweet potatoes	13	55	Decay, pitting, internal discoloration, hardcore when cooked
Tamarillos	3-4	37-40	Surface pitting, discoloration
Taro	10	50	Internal browning, decay
Tomatoes			
Ripe	7-10 [†]	45-50	Water soaking and softening, decay
Mature-green	13	55	Poor color when ripe, alternaria rot
Water convolvulus	10	50	Darkening of leaves and stems
Watermelons	4.5	40	Pitting, objectionable flavor

*Symptoms often become apparent only after removal to warm temperatures, as in marketing.

[†]See individual commodity sections in this Handbook.

Freezing Injury

The recommended storage temperatures for commodities that are not susceptible to chilling injury are as low as possible but slightly above the freezing point. Freezing injury occurs when ice crystals form in the tissues. Cultivars, locations, and growing conditions may affect the freezing point. To be on the safe side, the highest temperature at which freezing of a specific commodity may occur should be used as a guide for recommending the optimum storage temperature. More detailed discussion of freezing points and factors affecting them can be found in McColloch (1953), Whiteman (1957), and Parsons and Day (1970,1971). The most common symptom of freezing injury is a water-soaked appearance. Tissues injured by freezing generally lose rigidity and become mushy upon thawing.

The susceptibility of different fresh fruits and vegetables to freezing injury varies widely. Some commodities may be frozen and thawed a number of times with little or no injury, whereas others are permanently injured by even a slight freezing. All fruits and vegetables can be categorized into three groups based on their sensitivity to freezing: *most susceptible*—those that are likely to be injured by even one light freezing, *moderately susceptible*—those that will recover from one or two light freezing periods, and *least susceptible*—those that can be lightly frozen several times without serious damage. Table 2 shows the relative susceptibility of a number of fruits and vegetables to freezing injury.

Table 2. Susceptibility of fresh fruits and vegetables to freezing injury

Most susceptible	Moderately susceptible	Least susceptible
Apricots	Apples	Beets
Asparagus	Broccoli	Brussels sprouts
Avocados	Carrots	Cabbage, mature and savory
Bananas	Cauliflower	Dates
Beans, snap	Celery	Kale
Berries (except cranberries)	Cranberries	Kohlrabi
Cucumbers	Grapefruit	Parsnips
Eggplants	Grapes	Rutabagas
Lemons	Onion (dry)	Salsify
Lettuce	Oranges	Turnips
Limes	Parsley	
Okra	Pears	
Peaches	Peas	
Peppers, sweet	Radishes	
Plums	Spinach	
Potatoes	Squash, winter	
Squash, summer		
Sweet potatoes		
Tomatoes		

The freezing point of the commodity is no indication of the damage to be expected by freezing. For example, both tomatoes and parsnips have freezing points of -1.1 to -0.6 °C (30 to 31 °F), but parsnips can be frozen and thawed several times without apparent injury, whereas tomatoes are ruined after only one freezing. The severity of freezing injury is influenced by a combination of time and temperature. For example, apples that would be injured little by exposure to temperatures slightly below the freezing point for a few days would be severely injured by just a few hours of exposure to -7 to -10 °C (19 to 14 °F). The susceptibility to freezing injury is not necessarily similar for the same type of fruit or vegetable. For example, leafy lettuce is very susceptible to freezing injury, whereas some other leafy vegetables, such as kale and cabbage, can withstand several light freezing periods without serious injury.

When left undisturbed, most fruits and vegetables can usually be cooled one to several degrees below their freezing point before they actually freeze. This cooling without freezing is known as undercooling or supercooling. They may remain undercooled for several hours, but they will usually start to freeze immediately if jarred or moved. If permitted to warm above the freezing point, many commodities that have been undercooled may escape having ice crystals form in them. For example, potatoes, which are very sensitive to freezing damage, showed no freezing symptoms from having been undercooled for a short time to -4 °C (25 °F), about 3 °C (5 °F) below their freezing point, when they were carefully warmed after undercooling (Hruschka et al. 1961).

Plant tissues are very sensitive to bruising while frozen, and this sensitivity is another reason for leaving commodities undisturbed until they have warmed. Selecting a suitable thawing temperature involves a compromise. Fast thawing damages tissues, but very slow thawing such as at 0 to 1 °C (32 to 34 °F) allows ice to remain in the tissues too long and causes injury. Research on the rate of thawing has suggested that thawing at 4 °C (39 °F) causes the least damage for most commodities (Lutz 1936). Even though a number of fruits and vegetables are somewhat tolerant to freezing, commodities recovered from freezing often have shorter storage life and are more susceptible to invasion by microorganisms. For example, apples that recover from freezing are softer than normal fruit, and carrots that have been frozen are especially subject to decay. Therefore, it is best to avoid subjecting fresh produce to freezing temperatures in the first place.

References

- Bramlage, W.J., and S. Meir. 1990. Chilling injury of crops of temperate origin. *In* C.Y. Wang, ed., *Chilling Injury of Horticultural Crops*, pp. 37-49. CRC Press, Boca Raton, FL.
- Hruschka, H.W., R.V. Akeley, and E.H. Ralph. 1961. Seed potato productivity after cooling, supercooling or freezing. Marketing Research Report 507, U.S. Department of Agriculture, Washington, DC.
- Kosiyachinda, S., and R.E. Young. 1976. Chilling sensitivity of avocado fruit at different stages of the respiratory climacteric. *J. Amer. Soc. Hort. Sci.* 101:665-667.
- Lipton, W.J. 1978. Chilling injury of 'Honey Dew' muskmelons: symptoms and relation to

degree of ripeness at harvest. *HortScience* 13:45-46.

Lutz, J.M. 1936. The influence of rate of thawing on freezing injury of apples, potatoes and onions. *Proc. Amer. Soc. Hort. Sci.* 33:227-233.

Lyons, J.M. 1973. Chilling injury in plants. *Ann. Rev. Plant Physiol.* 24:445-466.

McColloch, L.P. 1953. Injuries from chilling and freezing. *In Plant Diseases, the Yearbook of Agriculture 1953*, pp. 826-830. U.S. Department of Agriculture, Washington, DC.

McColloch, L.P. 1962. *Alternaria* rot following chilling injury of acorn squashes. Marketing Research Report 518, U.S. Department of Agriculture, Washington, DC.

McColloch, L.P., and J.T. Worthington. 1952. Low temperature as a factor in the susceptibility of mature-green tomatoes to *Alternaria* rot. *Phytopathology* 42:425-427.

McColloch, L.P., J.N. Yeatman, and P. Loyd. 1966. Color changes and chilling injury of pink tomatoes held at various temperatures. Marketing Research Report 735, U.S. Department of Agriculture, Washington, DC.

Meir, S., S. Philosoph-Hadas, S. Lurie, et al. 1996. Reduction of chilling injury in stored avocado, grapefruit, and bell pepper by methyl jasmonate. *Can. J. Bot.* 74:870-874.

Parsons, C.S., and R.H. Day. 1970. Freezing injury of root crops: Beets, carrots, parsnips, radishes, and turnips. Marketing Research Report 866, U.S. Department of Agriculture, Washington, DC.

Parsons, C.S., and R.H. Day. 1971. Freezing injury to bell peppers. Marketing Research Report 895, U.S. Department of Agriculture, Washington, DC.

Paull, R.E. 1990. Chilling injury of crops of tropical and subtropical origin. *In C.Y. Wang, ed., Chilling Injury of Horticultural Crops*, pp. 17-36. CRC Press, Boca Raton, FL.

Raison, J.K., and G.R. Orr. 1990. Proposals for a better understanding of the molecular basis of chilling injury. *In C.Y. Wang, ed., Chilling Injury of Horticultural Crops*, pp. 145-164. CRC Press, Boca Raton, FL.

Ryall, A.L., and W.J. Lipton. 1979. Handling, Transportation, and Storage of Fruits and Vegetables, vol. 1, Vegetables and Melons, 2nd ed. AVI Publishing Company, Westport, CT.

Saltveit, M.E., and L.L. Morris. 1990. Overview on chilling injury of horticultural crops. *In C.Y. Wang, ed., Chilling Injury of Horticultural Crops*, pp. 3-15. CRC Press, Boca Raton, FL.

Wang, C.Y. 1982. Physiological and biochemical responses of plants to chilling stress. *HortScience* 17:173-186.

- Wang, C.Y. 1990. Chilling Injury of Horticultural Crops. CRC Press, Boca Raton, FL.
- Wang, C.Y. 1993. Approaches to reduce chilling injury of fruits and vegetables. Hort. Rev. 15:63-95.
- Wang, C.Y. 1994. Reduction of chilling injury by methyl jasmonate. Acta. Hort. 368:901-907.
- Wang, C.Y., and D.O. Adams. 1982. Chilling-induced ethylene production in cucumbers (*Cucumis sativus* L). Plant Physiol. 69:424-427.
- Whiteman, T.M. 1957. Freezing points of fruits, vegetables and florist stocks. Marketing Research Report 196, U.S. Department of Agriculture, Washington, DC.